

**Exam Three, MTH 205, Summer 2010**

Ayman Badawi

QUESTION 1. solve  $\frac{dy}{dx} = \frac{1-4x-2y}{2x+y}$

QUESTION 2. solve  $\frac{dy}{dx} = \frac{x^2 e^{(x^3+y^2)}}{y}$

**QUESTION 3.** Solve  $\frac{1}{x}y' + xy = \frac{x}{y^2}$

**QUESTION 4.** solve  $\frac{dy}{dx} = \frac{y^2 + x \ln(y) + x e^{x^2} + 3x}{-2yx - \frac{x^2}{2y} + \frac{\ln(y)}{y} + 10}$

QUESTION 5. Solve  $(2x + 2)y^{(2)} - 2y' = \frac{1}{x}$

**QUESTION 6.** Is there a solution to  $(2x + 2)y^{(2)} - 2y' = 0$ ,  $y(0) = 1$  and  $y'(-1) = 1$ ? If yes, find the solution. If not, does that contradict some of the results we studied? Explain.

**QUESTION 7.** A tank with capacity 20 liters contains 10 liters of water in which 20 grams of salt is dissolved. A mixture containing 1 gram of salt per liter is pumped into the tank at a rate 4 liters/min, the well-mixture solution is pumped out at rate 3 liters/min. Find the amount  $A(t)$  of salt in the tank at time  $t$ . What will be the amount of salt in the tank at the instant it overflows?

#### Faculty information

Ayman Badawi, Department of Mathematics & Statistics, American University of Sharjah, P.O. Box 26666, Sharjah, United Arab Emirates.

E-mail: [abadawi@aus.edu](mailto:abadawi@aus.edu), [www.ayman-badawi.com](http://www.ayman-badawi.com)

102

EXAM TWO, MTH 205, SPRING 008  
THIS IS THE REAL TEST

AYMAN BADAWI

ID NUMBER 2712

Name: Hashaa Rabbat

Score =

QUESTION 1. 20 points Solve  $y^{(2)} - 6y' + 9y = \frac{e^{3x}}{x^2} + e^{-3x}$

$y'' - 6y' + 9y = 0$

(homogeneous equation)

$y = e^{mx}$

$m^2 - 6m + 9 = (m-3)^2 = 0 \Rightarrow m = 3$

$y_c = c_1 e^{3x} + c_2 x e^{3x}$

$y_{p1} = A e^{-3x}$

$A 9e^{-3x} + A 18e^{-3x} + A 9e^{-3x} = e^{-3x}$

$A 36 e^{-3x} = e^{-3x} \Rightarrow A = 1/36$

$y_{p2} = f_1 e^{3x} + f_2 x e^{3x}$

$f_1' e^{3x} + f_2' x e^{3x} = 0$

$f_1' (3e^{3x}) + f_2' e^{3x} + 3x e^{3x} f_2' = \frac{e^{3x}}{x^2} \Rightarrow f_1' (3) + (1+3x)f_2' = \frac{1}{x^2}$

$b = \det \begin{bmatrix} e^{3x} & x e^{3x} \\ 3 & 1+3x \end{bmatrix} = e^{3x} + 3x e^{3x} - 3x e^{3x} = e^{3x}$

$f_1' = \det \begin{bmatrix} 0 & x e^{3x} \\ \frac{1}{x^2} & 1+3x \end{bmatrix} = \frac{-1 e^{3x}}{x} \cdot \frac{1}{e^{3x}} = -\frac{1}{x} \quad f_1 = \int f_1' = -\ln x$

$f_2' = \det \begin{bmatrix} e^{3x} & 0 \\ 3 & \frac{1}{x^2} \end{bmatrix} = \frac{e^{3x}}{x^2} \cdot \frac{1}{e^{3x}} = \frac{1}{x^2} \quad f_2 = \int \frac{1}{x^2} = -\frac{1}{x}$

$y_{p2} = -\ln x e^{3x} + (-\frac{1}{x}) x e^{3x}$

undetermined coefficient  
OK Good  
 $y_{p1} = \frac{e^{-3x}}{36}$   
(variant method)

Good

2

QUESTION 2. 20 points Solve  $y^{(2)} + \frac{1}{x}y' = \frac{1}{x^2}$

$$y'' + \frac{1}{x}y' = 0 \quad (\text{cauchy euler}).$$

$$y = x^m \quad y' = mx^{m-1} \quad y'' = m(m-1)x^{m-2}$$

$$m(m-1)x^{m-2} + \frac{1}{x}mx^{m-1} = 0$$

$$\Rightarrow m^2 - m + m = 0 = 0 = 0$$

$$y_c = c_1x^0 + c_2x^0 \ln x = c_1 + c_2 \ln x$$

$$y_p = f_1(x) + f_2(\ln x)$$

$$f_1' + f_2' \ln x = 0$$

$$0 + \frac{f_2'}{x} = \frac{1}{x^2} \Rightarrow f_2' = \frac{1}{x} \Rightarrow f_2 = \ln x$$

$$f_1' + \frac{\ln x}{x} = 0 \Rightarrow f_1' = -\frac{\ln x}{x}$$

$$f_1 = -\int \frac{\ln x}{x} = -\int u \, du \\ = -\frac{(\ln x)^2}{2}$$

$$\underline{y_g} = c_1 + c_2 \ln x - \frac{(\ln x)^2}{2} + \frac{(\ln x)^2}{2}$$

$$= c_1 + c_2 \ln x + \frac{(\ln x)^2}{2}$$

QUESTION 3. 20 points Solve  $y' + \cos(2x)y = \frac{\cos(2x)}{y} = \cos(2x)y^{-1}$

Bernoulli:

$$u = y^{1 - (-1)} = y^2$$

$$y = u^{1/2}$$

$$\frac{dy}{dx} = \frac{du}{du} \cdot \frac{du}{dx} = \frac{1}{2} u^{-1/2} u'$$

$$\rightarrow \frac{1}{2} u^{-1/2} u' + \cos(2x) u^{1/2} = \cos(2x) \left( \frac{1}{u^{1/2}} \right)$$

$$\frac{1}{2} u' + \cos(2x) u = \cos(2x)$$

$$u' + (2\cos(2x)) u = 2\cos(2x)$$

$$u = \frac{\int I \cdot F(x)}{I}$$

$$= \frac{\int 2\cos(2x) e^{\sin(2x)}}{e^{\sin(2x)}}$$

$$\frac{\int e^s du}{e^{\sin(2x)}} = \frac{e^{\sin(2x)} + c}{e^{\sin(2x)}} = u = \frac{1 + c}{e^{\sin(2x)}}$$

$$F(x) = 2\cos(2x)$$

$$I(x) = 2\cos(2x)$$

$$I = e^{\int 2\cos(2x) dx}$$

$$= e^{\sin(2x)}$$

$$= 1 + c e^{-\sin(2x)}$$

$$y = \sqrt{u} = \sqrt{\frac{e^{\sin(2x)}}{c + e^{\sin(2x)}}}$$

what!!



QUESTION 4. 20 points Solve  $y^{(2)} - \frac{2x}{x^2+1}y' + \frac{2}{x^2+1}y = \frac{2}{x^2-1}$  if  $y = x$  is a solution to the associated homogeneous system.

Reduction to first order: (homogeneous):

$$e^{-\int \frac{2x}{x^2+1} dx} = e^{\ln|x^2+1|} \quad \text{always +ve}$$

$$= |x^2+1| = (x^2+1)$$

$$y_2 = y_1 \int \frac{x^2+1}{y_1^2} dx = x \int \frac{x^2+1}{x^2} dx = x \int \left(1 + \frac{1}{x^2}\right) dx$$

$$= x \left(x - \frac{1}{x}\right) = \frac{x^2-1}{x}$$

$$y_c = c_1(x) + c_2(x^2-1)$$

$$y_p = f_1(x) + f_2(x^2-1)$$

$$f_1'(x) + f_2'(x^2-1) = 0$$

$$f_1' + f_2'(2x) = \frac{2}{x^2-1}$$

$$b = \det \begin{bmatrix} x & x^2-1 \\ 1 & 2x \end{bmatrix} = 2x^2 - x^2 + 1 = \underline{x^2+1}$$

$$f_1' = \det \begin{bmatrix} 0 & x^2-1 \\ \frac{2}{x^2-1} & 2x \end{bmatrix} = \frac{0 - 2(x^2-1)}{x^2-1} \cdot \frac{1}{x^2+1}$$

$$f_1 = \underline{-2 \tan^{-1}(x)}$$

$$f_2' = \det \begin{bmatrix} x & 0 \\ 1 & \frac{2}{x^2+1} \end{bmatrix} = \frac{2x}{x^2+1} = \frac{2x}{(x^2-1)(x^2+1)}$$

Good

$$u = x^2$$

$$f_2 = \int f_2' = \int \frac{1}{(u-1)(u+1)} du$$

$$f_2 = \int \frac{1}{2(u-1)} - \frac{1}{2(u+1)} du = \frac{1}{2} \ln|u-1| - \frac{1}{2} \ln|u+1|$$

$$= \underline{\frac{1}{2} \ln|x^2-1| - \frac{1}{2} \ln|x^2+1|}$$

+3

$$y_p = -2x \tan^{-1}x + (x^2-1) \left[ \frac{1}{2} \ln|x^2-1| - \frac{1}{2} \ln|x^2+1| \right]$$



**QUESTION 5. 20 points** A tank initially contains 20 gallons of Fresh WATER (i.e. when  $t = 0$ , amount of salt is zero). A mixture containing 0.5 bound of salt per gallon is poured into the tank at rate of 2 gallons per minute, while the mixture leaves the tank at rate 4 gallons per minute.

- Find the amount of salt in the tank at any time  $t$ .
- When will the tank be empty?
- Find the concentration of the salt in the tank at  $t = 9.5$  minutes.

DEPARTMENT OF MATHEMATICS & STATISTICS, AMERICAN UNIVERSITY OF SHARJAH, P.O. BOX 26666, SHARJAH, UNITED ARAB EMIRATES  
E-mail address: abadawi@aus.edu, www.ayman-badawi.com

$$V_0 = 20 \text{ gallons} = V(0) \quad A(0) = 0$$

$$C_{in} = 0.5 \text{ pound/gallon}$$

$$R_{in} = 2 \text{ gallons/min}$$

$$R_{out} = 4 \text{ gallons/min}$$

$$a) \quad \frac{dA}{dt} = R_{in} \cdot C_{in} - R_{out} C_{out}$$

$$V(t) = 20 + (R_{in} - R_{out})t = 20 - 2t$$

$$A' = 0.5(2) - \frac{A(t)}{20-2t} \cdot 4$$

$$C_{out} = \frac{A(t)}{V} = \frac{A(t)}{20-2t}$$

$$\Rightarrow A' + A(t) \left( \frac{4}{20-2t} \right) = 1$$

1st order linear:

$$I = e^{\int \frac{4}{20-2t} dt} = e^{\int \frac{2}{10-t} dt} = e^{-2 \ln|10-t|}$$

$$(if t > 10) \quad I = (10-t)^{-2}$$

$$A = \frac{\int (10-t)^{-2} dt}{(10-t)^{-2}} = \frac{+1(10-t)^{-1} + c}{(10-t)^{-2}} = (10-t) + c(10-t)^2 = A(t)$$

$$A(0) = 0 \Rightarrow 10 + c(10)^2 = 0 \Rightarrow c = \frac{-1}{10} = -0.1$$

$$A(t) = 10-t - 0.1(10-t)^2$$

$$b) \quad \text{when } V(t) = 0 \quad 0 = 20 - 2t \Rightarrow 20 = 2t \Rightarrow t = 10 \text{ minutes.}$$

At 10 minutes, the tank will be empty.

$$c) \quad A(9.5) = 10 - 9.5 - 0.1(10 - 9.5)^2 = 0.5 - 0.025 = 0.475 \text{ pounds.}$$